

VOLUMES OF REGULAR SEMISIMPLE HESSENBERG VARIETIES AND FACES OF GELFAND-ZETLIN POLYTOPES

MIKIYA MASUDA (OSAKA CITY UNIVERSITY)

If X is a nonsingular projective variety of complex dimension d together with an embedding $X \hookrightarrow \mathbb{P}^N$, then the volume of the embedding is defined by

$$\text{vol}(X \hookrightarrow \mathbb{P}^N) := \frac{1}{d!} \deg(X \hookrightarrow \mathbb{P}^N) = \frac{1}{d!} \int_X c_1(L)^d$$

where L is the very ample line bundle on X associated to the embedding. If X is a flag variety $\text{Flags}(\mathbb{C}^n)$ together with a Plücker embedding, then its volume is known to agree with the volume of the associated Gelfand-Zetlin polytope.

Regular semisimple Hessenberg varieties are nonsingular subvarieties of $\text{Flags}(\mathbb{C}^n)$, so the Plücker embedding of $\text{Flags}(\mathbb{C}^n)$ induces their embeddings to \mathbb{P}^N . In this talk, we discuss how to represent their volumes in terms of faces of the Gelfand-Zetlin polytope.

This is joint work with Megumi Harada, Tatsuya Horiguchi, and Seonjeong Park.

REFERENCES

- [1] H. Abe, L. Dedieu, F. Galetto, and M. Harada, *Geometry of Hessenberg varieties with applications to Newton-Okounkov bodies*, *Selecta Math.* (to appear); arXiv:1612.08831.
- [2] T. Abe, T. Horiguchi, M. Masuda, S. Murai, and T. Sato, *Hessenberg varieties and hyperplane arrangements*, arXiv:1611.00269.
- [3] V. Kiritchenko, E. Smirnov, and V. Timorin, *Schubert calculus and Gelfand-Zetlin polytopes*, *Russian Math. Surveys*, 2012, 67 (4) 685-719; arXiv:1101.0278.